

# **Appendix AAE**

**Reynolds, R.T., R.T. Graham, and D.A. Boyce. 2006a. An Ecosystem-Based Conservation Strategy for the Northern Goshawk. Studies in Avian Biology. In press.**

In Press

Document cover Sheet—Studies in Avian Biology 31

Printing order:

Title: AN ECOSYSTEM-BASED CONSERVATION STRATEGY FOR THE NORTHERN GOSHAWK

RH: Goshawk Conservation Strategy—*Reynolds* et al.

Number of text pages: 44

Number of figures: 4

Number of tables: 1

Author to receive correspondence:

Richard T. Reynolds  
USDA Forest Service  
Rocky Mountain Station  
2150 Centre Ave., Suite 350, Bldg. A  
Fort Collins, CO 80526-1891  
e-mail: [rreynolds@fs.fed.us](mailto:rreynolds@fs.fed.us)

Carl D. Marti, Editor  
Studies in Avian Biology  
Raptor Research Center  
Boise State University  
Boise, ID 83725 USA  
Tel: 208-426-8996  
Fax: 208-426-3117  
Email: [cmarti@boisestate.edu](mailto:cmarti@boisestate.edu)

# AN ECOSYSTEM-BASED CONSERVATION STRATEGY FOR THE NORTHERN GOSHAWK

RICHARD T. REYNOLDS, RUSSELL T. GRAHAM, AND DOUGLAS A. BOYCE, JR

*Abstract.* The Northern Goshawk (*Accipiter gentilis*) is a large forest-dwelling raptor whose viability is in question because of habitat changes resulting from tree cutting, fire exclusion, and livestock grazing. We describe an approach for developing a goshawk habitat conservation strategy, first used in the southwestern United States in 1992, that can be applied throughout the range of the species. The strategy described sets of desired habitats based on existing knowledge of the life history and habitats of goshawks, the life histories and habitats of their prey, and the ecology of overstory and understory vegetation in forests occupied by goshawks. These habitats included components such as overstory and understory compositions and structures, snags, logs, woody debris, openings, and size (area) and arrangement of plant aggregations. The strategy incorporated the dynamic nature of forest ecosystems by developing desired landscapes consisting of temporally shifting mosaics of vegetation structural stages that comprised the habitats of goshawks and their prey. This multi-species, ecosystem-based strategy will benefit goshawks because their populations are limited by food and habitat and because the desired landscape will contain goshawk and their prey habitats through time. The approach used in this conservation strategy should be appropriate for other forests occupied by goshawks. However, because the species of prey, and the composition, structure, and dynamics of the vegetation vary among forest types, the approach is likely to result in unique desired habitats and landscapes as well as forest management prescriptions to develop them.

*Key Words:* *Accipiter gentilis*, conservation strategy, food webs, forest management, habitat, landscapes, prey, structural stage.

Considerable effort has been directed towards developing conservation strategies that protect forest species. Many conservation strategies prompted by recovery goals in the Endangered Species Act are autecological, spatially and temporally limited, and typically use habitat reserve designs (Everett and Lehmkuhl 1996, but see DellaSala et al. 1996, MacCracken 1996, Noss 1996, and Everett and Lehmkuhl 1997 for discussions on the merits of reserves). These strategies often fail to recognize important ecological relationships and linkages that support a species (e.g., food webs) and they often view habitats as static. Although reserves may protect species that are sensitive to human activities, their very design shifts resource extraction pressures to unprotected areas, which may diminish the ecological values of reserves by limiting dispersal (gene flow) of focal species among reserves (Suzuki 2003). Conservation strategies that address all stages of a species' life history, the physical and biological factors that limit its populations, the members of its ecological community, and the spatial and temporal dynamics of the ecosystems it occupies, should be robust to failure. Implementing such strategies in landscapes increases the probability of sustaining whole ecosystems on which a species may depend, and eliminates the difficult tasks of specifying the sizes, numbers, dispersion, and connectivity of reserves or protected areas needed to sustain a species.

Apex predators, because they are often sensitive to changes in their habitats (Belovsky 1987, Melián and Bascompte 2002), are prime candidates for conservation strategies. Population viability of the Northern Goshawk (*Accipiter gentilis*), an apex predator that occurs primarily in forests and woodlands throughout the Holarctic (Squires and Reynolds 1997), is in question because of habitat changes resulting from tree cutting, fire exclusion, and livestock grazing (Herron et al. 1985, Crocker-Bedford 1990, Reynolds et al. 1992, Widén 1997, but see Kennedy 1998). As a result, goshawks have been the object of considerable litigation and the species was

considered for listing under the Endangered Species Act (Boyce et al., *this volume*). To protect the habitats of goshawks, conservation strategies were developed for three forest types in the southwestern United States in 1992 (Reynolds et al. 1992). These southwestern goshawk conservation strategies (SWGS) accounted for the requisite resources (habitat and food) and ecological relationships (competition, predation, and disease) of goshawks and their prey. Further, because forests change through the dynamic processes of plant establishment, growth, succession, and natural and anthropogenic disturbances, the SWGS identified and incorporated the spatial and temporal scales encompassing these dynamics. The SWGS described sets of desired forest conditions that included habitat components such as tree species composition, structure, landscape pattern, snags, woody debris, tree sizes and densities, and the sizes, ages, and arrangement of tree groups. To account for forest dynamics, the desired forest conditions consisted of temporally shifting mosaics of vegetation structural stages intended to sustain the habitats of both goshawks and their prey in large landscapes for centuries.

The SWGS was incorporated into all USDA Forest Service Southwestern National Forest management plans in 1996 (Boyce et al., *this volume*, USDA Forest Service 1996). Shortly thereafter, the SWGS was reviewed by animal and forest scientists (Braun et al. 1996, Squires et al. 1998, Long and Smith 2000, Peck 2000, Beier and Maschinski 2003, Andersen et al. 2004). Here we review the approach, components, and processes used in the SWGS, particularly those applicable to southwestern ponderosa pine (*Pinus ponderosa*) forests, not only to correct misunderstandings evident in some of the reviews, but to demonstrate how the process can be used to develop similar conservation strategies in other forests. We conclude with a discussion of problems that may hinder tests of the effectiveness of the SWGS for sustaining goshawks and identify some unintended, additional values resulting from implementation of the SWGS.

## ESSENTIAL INFORMATION

Information on the life history, ecology, and habitat of the goshawk, the biological and physical factors (food, habitat, predators, competitors, disease, and weather) that potentially limit goshawk populations, the life histories and populations of important goshawk prey species, and the ecology (e.g., composition, structure, pattern, and dynamics) of a forest ecosystem, is essential for developing desired forest conditions in this ecosystem-based conservation strategy.

## GOSHAWK LIFE HISTORY

Goshawks are relatively long-lived, solitary breeders with large home ranges, and that breed in a broad range of forest and woodland types (Squires and Reynolds 1997) where they feed on a variety of birds and mammals (Reynolds and Meslow 1984, Boal and Mannan 1994, Reynolds et al. 1994). Goshawks exhibit high levels of year-to-year fidelity to breeding territories and to mates (Doyle and Smith 1994, Woodbridge and Detrich 1994, Squires and Ruggiero 1995, Reynolds et al. 1994), and often lay eggs in numerous alternate nests within their territories (Reynolds et al. 1992, Woodbridge and Detrich 1994, Reynolds and Joy, *this volume*). Studies have shown that where forests have suitable structures for nests and hunting, and where food is abundant, goshawks are more abundant, breed more often, have heavier body masses, and smaller home ranges (McGowan 1975, Bednarek et al. 1975, Sollien 1979, Lindén and Wikman 1980, Cramp and Simmons 1980, Sulkava et al. 1994, Reynolds et al. *this volume*).

## GOSHAWK LIMITING FACTORS

A fundamental step in developing conservation strategies is to identify the environmental factors that limit a goshawk population's ability to grow. These factors typically affect goshawk birth, death, emigration, and immigration rates. Sources of information for these factors include the published literature, unpublished reports, and expert opinion. Information on factors that may

limit goshawk populations is often scarce or absent. In these cases, information on how factors influence other raptor populations may offer indications on how they might influence goshawks. A recent review of the goshawk and other raptor literature identified factors that may limit goshawk populations—the abundance and availability of habitats and foods, the types and abundances of predators and competitors, diseases, and weather (Reynolds et al., *this volume*). The review also showed that in studies of goshawk breeding density and reproduction, the availabilities of nest sites, foods, and suitable foraging sites appeared to be the most common factors affecting goshawk populations, and that predation, competition, disease, and weather would be less likely to affect goshawks negatively if foods and habitats were not limiting (Reynolds et al., *this volume*). For example, when prey are abundant, competition for food might be reduced, food stress would less likely predispose goshawks to disease, weather effects on prey availability might be reduced, and, when high quality nest sites are available, predation at goshawk nests might be reduced (Reynolds et al. 1992). The conservation problem was then to identify and develop the habitats of sufficient quality to support goshawks and their prey populations. The variation among habitats in the composite availabilities of nest sites, foraging sites, foods, escape cover, and abundances of predators and competitors determines habitat quality. The approach used in SWGS assumes that if high quality habitats are available in landscapes then the above limiting factors would less likely constrain the growth of goshawk populations.

#### GOSHAWK HABITAT

NORTH American goshawks nest and hunt in a wide variety of many forest and woodland types within their geographic range (Squires and Reynolds 1997). Based on the use of space around goshawk nests by adults and fledglings, the SWGS identified three components of the

breeding home range: the nest area (approximately 12 ha), the post-fledging family area (PFA; approximately 170 ha exclusive of nest area) surrounding the nest area, and the foraging area (approximately 2,190 ha exclusive of PFA) surrounding the PFA (Reynolds et al. 1992). We know more about the composition and structure of vegetation in nest areas than in the other areas because of their small size, readily defined boundaries, and the numerous studies that described nest site and nest area vegetation. Forests within nest areas provide protected nest, roost, and prey handling sites (Reynolds et al. 1982). Little foraging occurs within nest areas (Schnell 1958) and nest area sizes and shapes can vary by landform, forest setting, and method used to quantify them (Reynolds 1983, Kennedy 1989, Kennedy 1990, Boal et al. 2003). Goshawk nest areas typically have relatively high densities of large trees, inherent to the forest type and biophysical setting, relatively high canopy cover, open understories, and are typically on shallow slopes or in drainages protected by such slopes (Squires and Reynolds 1997). While most nest areas are embedded within extensive forests or woodlands, some goshawk individuals and populations nest in small patches of trees within open shrub, tundra, or riparian habitats (Bond 1940, White et al. 1965, Swem and Adams 1992, Younk and Bechard 1994a, b). Despite the disparate species compositions of forest types used by breeding goshawks, the structure of forests within nest areas is surprisingly consistent suggesting that structure is more important than species composition in their choice of nest habitat.

The PFA, defined in the SWGS as the adult female core area including the nest (Kennedy 1989), is used by the adult female for foraging and by her fledglings during the post-fledging dependency period (Reynolds et al. 1992). Because PFAs are larger than nest areas, they typically include a wider diversity of forest conditions (species compositions, age classes, openings, and landforms). Because goshawk fledglings spend much of the post-fledging dependency period near



the center of a PFA where they may require additional hiding cover from predators, the desired PFA habitat condition is a transition from the denser forests in nest areas to more open foraging habitat in the outer portions (Reynolds et al. 1992).

The foraging area surrounds the PFA and comprises the remainder of the home range of breeding goshawks (Reynolds et al. 1992). The foraging area is used by adult goshawks for hunting, and, like the PFA, should comprise suitably structured foraging habitat and a mix of prey habitats (Reynolds et al. 1992). A number of radio-telemetry studies determined the use of habitats by goshawks (Kenward et al. 1981, Widén 1985, Kenward and Widén 1989, Bright-Smith and Mannan 1994, Hargis et al. 1994, Squires and Ruggiero 1995, Beier and Drennan 1997, Good 1998, Drennan and Beier 2003), but their elusive behavior and rapid movements through large home ranges make goshawks difficult to observe and to unequivocally determine whether or not they were actually hunting in the habitats they were detected using. Nonetheless, these studies suggested that breeding goshawks hunted primarily in mature and old forests, but that they also hunted in a variety of other forest age classes, structures, and compositions, and into openings and along forest edges (see White et al. 1965, Widén 1989, Bright-Smith and Mannan 1994, Hargis et al. 1994, Younk and Bechard 1994a, Younk and Bechard 1994b, Bosakowski et al. 1999, Daw and DeStefano 2001). The diversity of habitats used by hunting goshawks often expands during winter when many juveniles and some adults move to lower elevation woodland and shrub communities (Reynolds et al. 1994, Squires and Ruggiero 1995, Stephens 2001, Sonsthagen 2002). Whether these goshawks leave their forest habitats in response to reduced food availability or weather changes is unknown. This diversity of habitat use by goshawks is often reflected in their diets; goshawks eat birds and mammals that occur in mature forests, but frequently eat species whose main habitats are in open forests (e.g., ponderosa

pine), along forest edges, and in openings (e.g., meadows) (Reynolds and Meslow 1984, Widén 1989, Boal and Mannan 1994, Daw and DeStefano 2001). Nonetheless, at least within forest situations, goshawks spend much of their time in areas with large trees (Bright-Smith and Mannan 1994, Hargis et al. 1994), areas with high crown–base heights (open understories), allowing goshawks to fly beneath the forest canopy. Older forests also contain abundant tree perches from which goshawks search for prey, and are the prime habitat of many goshawk prey species (Reynolds et al. 1992).

#### GOSHAWK PREY

Goshawks feed on birds and small mammals (Squires and Reynolds 1997), and the composition of a local goshawk diet depends on the composition of the bird and mammal fauna in a particular forest, the relative abundances and availabilities of the species that goshawks are able to capture, and the dietary preferences of the goshawks. Goshawk diets comprise a limited range of prey sizes (Storer 1966, Snyder and Wiley 1976, Reynolds and Meslow 1984, Bosakowski et al. 1992). The upper prey-size limit appears to be determined by the goshawk's ability to kill with a minimum risk of injury to itself, and the lower size limit is likely determined by a goshawk's ability to capture smaller prey. Small prey are more maneuverable and escape goshawks more readily and return less energy per capture than larger prey (Reynolds 1972, Andersson and Norberg 1981, Temeles 1985). These limits result in goshawk diets composed of robin-to-grouse-sized birds and chipmunk-to-hare-sized mammals (Reynolds et al. 1992).

Goshawks are morphologically and behaviorally suited to hunt in forests. Both their maneuverability for capturing agile prey, provided by short wings and long tail, and their short-perch-short-flight foraging tactic (Kenward 1982), are suited for environments where flight and vision is impaired by tall, dense vegetation (Reynolds et al. 1992). Because of these adaptations it

is often assumed that goshawks are limited to old-growth forests and that habitat availability is the main factor limiting goshawk populations. However, even within the forests, goshawk reproduction and survival can be highly variable among years (Reich et al. 2004, Keane et al., *this volume*, Reynolds and Joy, *this volume*), and this variation has been associated with inter-annual variations in prey abundance ( McGowan 1975, Lindén and Wikman 1980, Doyle and Smith 1994, Selås 1997, Keane 1999, Salafsky 2004). Furthermore, Widén (1989) reported higher breeding densities in areas richer in foods, and Bednarek et al. (1975) reported extremely high goshawk breeding densities in areas with only 12–15 % of woodland but very rich in food. Widén (1989) suggested that goshawks are more often limited by food than by nesting habitat.

#### GEOGRAPHIC AND ANNUAL VARIATION IN DIETS

Goshawk diets differ among forest types, among regions, and both seasonally and annually. Reynolds and Meslow (1984), Kennedy (1991), and Boal and Mannan (1994) reported between 14 and 37 different prey species in goshawk diets in a variety of western American conifer forests, while in eastern American deciduous forests, 23 different prey species were reported (Bosakowski and Smith 1992, Bosakowski et al. 1992). Much of the among-forest and regional differences in diets disappears, however, when prey are grouped at the genus level because prey species are often regionally replaced by congeners. For example, red squirrels (*Tamiasciurus hudsonicus*) in western Oregon are replaced by Douglas squirrels (*Tamiasciurus douglasi*) in eastern Oregon and Nuttall's cottontail (*Sylvilagus nuttalli*) in western North America is replaced by the eastern cottontail (*Sylvilagus floridanus*) in eastern North American (Hall 1981). Due to such replacements, goshawk diets can be generalized to include rabbits, tree squirrels, ground squirrels, woodpeckers, jays, thrushes, doves, pigeons, and grouse. However, goshawks frequently supplement these prey with as many as 20 other incidental bird and

mammal species (Schnell 1958, Reynolds and Meslow 1984).

Annual variation in local goshawk diets may stem from annual variation in prey abundances associated with eruptive or inter-annual fluctuations in species such as snowshoe hare (*Lepus americanus*), red squirrel, and grouse (e.g., *Dendragapus obscurus*) (McGowan 1975, Doyle and Smith 1994). Although little winter goshawk diet information is available, diets are likely to vary seasonally due to habitat differences among prey, differential sampling of habitats by foraging goshawks, and the timing of estivation, hibernation, or migration of some prey. The abundance of non-migratory prey (tree squirrels, hares, grouse, woodpeckers) during winter may affect whether goshawks stay on breeding territories or move to non-forest habitats in winter.

#### DETERMINING DIETS

Because the SWGS approach for developing conservation strategies requires the identification of a suite of important goshawk prey in a focal forest type, we review methods for estimating goshawk diets and a process that can be used to reduce a complete list of prey in a forest type to a reduced list of important prey. Most of our understanding of goshawk diets comes from the breeding period when prey is delivered to nests by adults. Breeding season diets have been estimated with several methods, each with a characteristic bias. A prey-remains method takes advantage of the fact that goshawks regurgitate pellets and pluck feathers and fur from prey in their nest areas (Reynolds and Meslow 1984, Martin 1987). A bias associated with this method is inaccurate counts of individuals or species due to species-specific differences in detectability of remains when they are being collected (Reynolds and Meslow 1984, Bielefeldt et al. 1992). A direct observation method involves identifying and counting prey delivered to nests from adjacent blinds or with cameras (Schnell 1958, Boal and Mannan 1994, Grønnesby and Nygård

2000). Problems with direct observations are that the number of nests that can be observed is typically limited and difficulty of identifying prey whose diagnostic parts (feathers, fur) have been removed by the goshawks. Schnell (1958) identified 14 prey species from observations at a single nest in California, whereas Reynolds and Meslow (1984) identified 37 different species from prey remains collected at 58 goshawk nests in Oregon. Diet studies that combine these two methods are likely to result in more precise estimates of goshawk diets, but neither method accounts for prey eaten away from nests (Lewis et al. 2004).

What little we know about non-breeding season diets comes mostly from radio-telemetry study of wintering goshawks (Kenward 1979, Widén 1987, Stephens 2001, Drennan and Beier 2003, Tornberg and Colpaert 2001). Diets of goshawks that remain in forests during winter are not likely to differ greatly from the breeding diets, except prey that hibernate or migrate will be missing, and diets of goshawks that move to open habitats are more likely to include non-forest prey. Of course, diets should be determined from an adequate sample of goshawks within a forest type to reduce sampling error (i.e., a goshawk taking aquatic birds from a lake), and should be determined over an adequate number of years to detect inter-annual fluctuations in prey species.

#### SUITES OF IMPORTANT PREY

A reduction in a complete list of goshawk prey in a forest to a subset of important goshawk prey may be necessary because some species are taken only incidentally and their inclusion might dilute the forest habitats needed by more commonly captured prey. Goshawk diets are rarely dominated by a few species. However, typically less than half the species eaten contribute to more than a majority of individuals consumed. In California, six of a total 14 prey species contributed about 80% of the numbers of prey in the diet of a single goshawk pair (Schnell 1958), 18 of 37 species contributed 85% of prey in a large sample of Oregon nests

(Reynolds and Meslow 1984), and 11 of 18 species contributed 67% of prey in Arizona (Boal and Mannan 1994). Also, rarely does a single prey species contribute more than 30% of total numbers of prey in a diet; in fact, most prey species contributes less than 5% of the total. If a threshold for identifying a suite of important prey was chosen to include all species contributing more than 2% of individuals in a goshawk diet, then the suite would include eight prey species (57 % of total species) in Schnell's (1958) California study, 18 species (49%) in Reynolds and Meslow's (1984) Oregon study, and 11 species (61%) in Boal and Mannan's (1994) Arizona study.

However, because larger prey contribute more food biomass to the energy budget of goshawks, they can be more important than small prey even though small prey are eaten more often. Using the above 2% threshold in Table 1 excludes three large species (e.g., Belding's ground squirrel (*Citellus beldingi*), mountain cottontail (*Sylvilagus nuttalli*), Ruffed Grouse (*Bonasa umbellus*) that probably should be included in a suite of important prey because of their body mass. In Table 1, thresholds lower than two individuals per species may include too many incidental prey. Alternatively, including too few species may result in an insufficiently diverse and abundant food resource to sustain goshawks through poor food years. Other information, such as a comparison of the abundance of a marginally important prey species in unmanaged forests to its abundance (and frequency in goshawk diets) in managed forests, may help decide on whether or not to include marginal species in the suite of important prey. Finally, it must be pointed out that the diversity of habitats provided for the suite of 14 prey species in southwestern forests also provided habitats for many incidental prey species as well as non-prey species (Reynolds et al. 1992).

#### PREY HABITATS

After identifying a suite of important prey, the distributions, life histories, densities, and habitats of the prey can be assessed in the literature (e.g., journals, theses, and reports) and by expert opinion (Reynolds et al. 1992). Much information on the ecology and habitats of a variety of goshawk prey is already in Reynolds et al. (1992) and Drennan et al. (*this volume*). Often, information on the ecologies, habitat relations, and foods of prey species within a certain forest type is limited. In these cases, information from the same or a similar forest type in adjacent regions could be used. Limitations of these kinds of data include: (1) incomplete information on a species' life histories, population ecologies, and how these vary among forest types; (2) uncertainty about relationships between a species' demography and habitat conditions; (3) difficulties distinguishing a species' habitat use from its habitat preference; and (4) the appropriateness of using studies designed to investigate other questions (Morin 1981, VanHorne 1983, White and Garrott 1990).

#### FOREST HABITAT ELEMENTS OF PREY

Once the life histories, habitats, and foods of important prey are assessed, a list of forest habitat elements (FHE), including items such as vegetation structural stages, size of openings, edges, understory and overstory compositions and structures, woody debris, snags, nesting and feeding substrates, and interspersions of forest age classes, for each prey species can be developed. This process can be facilitated with matrices that display the frequencies of the relative importance (high, medium, low, or none) of FHEs for each prey species (see Table 6 in Reynolds et al. 1992). An overall relative importance of FHEs for the suite of prey can be estimated by summing the levels of importance of each FHE across species (Table 6 in Reynolds et al. 1992). Such assessments for the suite of goshawk prey in southwestern forests resulted in an understanding of the importance of sustaining large amounts of mid-aged to old forests

dispersed at a fine scale within landscapes (Reynolds et al. 1992, Long and Smith 2000).

## FOREST ECOLOGY

Forests, and by extension forest habitats, are dynamic ecosystems that undergo change through plant growth and succession and periodic natural and anthropogenic disturbances such as wind, fire, insects, and vegetation management. Each of these factors changes the composition, structure, and pattern of plant communities, which in turn have short- and long-term effects on wildlife habitats. Thus, describing and managing forest habitats for plants and animals in the goshawk food web requires an understanding of forest dynamics as well as the habitat relationships of the plants and animals. Here we identify sources of essential information on how to develop and sustain desired forest conditions through management, how to identify limits or constraints on such parameters as maximum tree sizes and longevity, sizes of plant aggregations and tree densities, and the species composition, structure, and landscape pattern of desired landscapes. Some important processes that occur during forest development include plant establishment, development, senescence, competitive exclusion, biomass accumulation, canopy gap initiation, understory reinitiation, maturation, decadence development, and mortality (Franklin et al. 2002). Each of these processes, which may vary among forest types, is often integrated into potential vegetation classifications. Moreover, these classifications provide estimates of forest productivity, vegetation development rates, plant occurrence and position (e.g., canopy layer), life form (e.g., grass, forb, or shrub), their roles in plant succession (e.g., early, mid-, or late seral), and include physical and biological components such as climate, soil, geology, and vegetation (Daubenmire and Daubenmire 1968, Cooper et al. 1991, Hann et al. 1997). These classification systems can also be integrated with known fire relations (Bradley et al. 1992, Agee 1993, Hann et al. 1997, Graham et al. 1999, Kaufmann et al. 2000) and are



compatible with efforts for defining and mapping fire regime condition classes for forests (Schmidt et al. 2002). Sources of data on current forest conditions include Forest Inventory and Analysis and Geospatial Analysis Processes (USGS National Gap Analysis Program 1995, O'Brien 2002).

## SYNTHESIS OF COMPONENTS

Once information on goshawks, their prey, and forest ecology is assembled, it is synthesized into desired sustaining mixes of goshawk and prey habitats. The SWGS used a vegetation structural stage (VSS) classification to describe forest development (Fig. 1). VSS is an integrative approach that combines vegetation growth and maturation into generalized descriptions of forest conditions from young to old vegetation complexes (Thomas et al. 1979, Verner and Boss 1980, Oliver and Larson 1990, Franklin et al. 2002, and see Table 7 and Appendix 5 in Reynolds et al. 1992). The FHEs were incorporated with VSS in a generalized landscape that included an abundant and dispersed large-tree component (large live trees, large snags, and large logs), groups (<0.2 ha) of trees with interlocking crowns, small openings (Fig. 1) Fifaround tree groups with a well-developed grass/forb/shrub vegetation, and a high level of interspersions (intermixing) of VSS in small (< 0.2 ha) patches of trees (Reynolds et al. 1992, Long and Smith 2000) (Fig. 2). Groups of trees with interlocking crowns allow the tasseled-eared squirrel (*Sciurus aberti*) and red squirrel (*Tamiasciurus hudsonicus*) to move among tree crowns, a critical habitat element especially around their nests (Reynolds et al. 1992 and citations therein, Dodd et al. 2003). Because mycorrhizal fungi are an important food for squirrels, and because the fungi are more abundant in mid-aged forests, an interspersions of mature and old VSS groups with mid-aged VSS groups benefits these squirrels. Small (Fig. 1) openings containing grass/forb/shrubs around tree groups are habitat for prey such as rabbits, ground squirrels, and

grouse that require openings for feeding or brood rearing. These openings should be maintained as natural openings because they are occupied by roots of the grouped trees (Pearson 1950), facilitating nutrient uptake and vigorous tree growth. Openings (e.g., <1.5 ha), because they are occupied by certain prey, also offer hunting opportunities for goshawks (Reynolds et al. 1992).

#### FOREST SETTING

An integration of information on the autecology and synecology of forest vegetation is essential for developing and sustaining goshawk and prey habitats (Fig. 1). A wealth of information on forest development can provide guidance for the development of the desired habitats. This information includes, but is not limited to, vegetation classifications, forest vegetation simulations, fire histories, natural-area descriptions, and wild-land, fuel-management strategies (e.g., Haig et al. 1941, Pearson 1950, Daubenmire and Daubenmire 1968, White 1985, Fulé et al. 1997, Reinhardt and Crookston 2003, Graham et al. 2004). Such information provides verification that can increase the likelihood that the desired goshawk and prey habitats in a particular forest type can be attained and sustained.

Sustaining the desired landscape mix of goshawk and prey habitats over space and time requires the incorporation of the spatial and temporal dynamics of forest vegetation. Vegetation dynamics, including the establishment, development, senescence, and its composition, structure, and pattern, can be estimated and modeled (Oliver and Larson 1990, Reynolds et al. 1992, Franklin et al. 2002, Reinhardt and Crookston 2003). For example, sustaining the maximum amount of mature and old VSS in southwestern forests for goshawks and their prey was best achieved with about 10% of landscape in VSS 1 (grass/forb/shrub), 10% in VSS 2 (seedling/sapling), 20% in VSS 3 (young forest), 20% in VSS 4 (mid-aged forest), 20% in VSS 5 (mature forest), and 20% in VSS 6 (old forest) (Reynolds et al. 1992). These proportions reflect

forest development from cohort establishment through canopy closure to old forests. However, classification systems that depict forest development over 1,000 yr tend to display greater proportions of a forest in the mature and old classes than classification systems depicting forest development through periods <300 yr. For example, Franklin et al. (2002) showed over 70% of the forest occurring in structural stages greater than 800 yr, as did Spies and Franklin (1996). Integrating a VSS distribution with goshawk habitats (nest area, PFA, foraging area) and tree-group metrics favoring the suite of southwestern prey, resulted in desired landscapes comprised of shifting mosaics of VSS through time and space (Reynolds et al. 1992, Long and Smith 2000).

The above syntheses produced desired forest conditions containing both goshawk and prey habitats. Remarkably, the desired conditions developed in the SWGS approximated the composition, structure, and landscape pattern existing in southwestern ponderosa pine forests before fundamental changes in natural disturbance regimes (Pearson 1950, White 1985, Fulé et al. 1997, Long and Smith 2000) (Fig. 2). Of course, it is important that the animal habitat relations and vegetation ecology information used to develop ecosystem-based conservation strategies be internally consistent as well as consistent with the current knowledge (Guldin et al. 2003).

#### IMPLEMENTATION OF THE GOSHAWK STRATEGY

Once the desired compositions, structures, and mixes of goshawk and prey habitats are described, management actions can be developed and implemented through appropriate planning processes. The SWGS recommended that goshawk breeding habitat be partitioned into nest areas, PFAs, and foraging areas, and because the movements of breeding goshawks are energetically limited to some finite space around their nests, that these home range components be approximately centered on the nest. Goshawk conservation strategies can be implemented at a

variety of spatial scales depending on management objectives. For example, implementation at the goshawk home range scale is appropriate for developing and protecting habitats in known territories. If the intent is to provide habitat for undiscovered goshawks or for an expansion of a goshawk population, the scale must be larger, e.g., a national forest or ecoregion (Reynolds et al. 1992, Graham et al. 1999). Implementing the strategy in entire landscapes accommodates seasonal, annual, and geographic variation in goshawk home range sizes (Hargis et al. 1994, Boal et al. 2003), and eliminates the need to specify the number of breeding territories, their juxtaposition and connectivity to sustain goshawk populations.

Specific management actions and the intensity that they are applied should be contingent on the differences between the existing conditions of a focal area and the desired conditions. If differences are great (e.g., no old-forest structure), centuries may be needed to develop the desired conditions. For example, more than 200 yr are required to develop old-forest structure in southwestern ponderosa pine forests (Reynolds et al. 1992), and over 1,200 yr are required to develop all of the structural stages found in northwestern Douglas-fir forests (Franklin et al. 2002). The capability of forests to produce the desired conditions can vary among sites depending on factors such as soils, slope, exposure, elevation (Daubenmire and Daubenmire 1968, Wykoff and Monserud 1988, Basset et al. 1994). Thus, differing growth potentials require that site-specific desired conditions be matched to a site's capabilities. Not all sites within a landscape can, nor should they have, the same exact conditions.

The Kaibab National Forest in Arizona began implementing the SWGS in ponderosa pine forests in 1993. Fig. 3 displays one such implementation (left portion of photo) adjacent to 12–16 ha seed-tree cuts (center, lower right), a forest treatment in which a few trees are retained as seed sources, and a natural area (top center) that had recent low intensity surface fires and little tree

cutting. Note the similarities in the aggregation of ponderosa pine trees and surrounding openings in the implementation area and the natural area. A lesson learned from multiple implementations is to avoid removing trees from within groups (especially in mid-aged, mature, and old VSS). Thinning groups often eliminates the interlocking of tree crowns, critical habitat for tree squirrels (Dodd et al. 2003). Rather, when tree cutting is needed to create or sustain the desired conditions, an entire group of trees should be regenerated as opposed to thinning within a group. The desired within-group structures in both mature and old VSS, could be developed with appropriate forest treatments (e.g., thinning or prescribed fire) in the younger age classes (e.g., seedling/sapling, young forests, and mid-aged forests) (Reynolds et al. 1992).

#### EVALUATION OF IMPLEMENTATION

Squires et al. (1998) suggested that the SWGS be tested before large scale implementation. Testing is needed to determine if management actions successfully moved existing forest conditions toward the desired conditions and if the actions had the desired effects on goshawks and their prey. One such test is to compare goshawk reproduction and survival in forests that are in or near the desired conditions to those in contrasting forests (paired landscape approach). Such comparisons, however, could be confounded by ecological differences (e.g., soil types) in the areas being compared. Another approach is to monitor the effects of implementation on the same sample of goshawk territories over time (before and after treatment design). However, depending on the degree of difference between existing and desired forest conditions, and because forest annual treatments are typically small relative to goshawk home ranges, achieving the desired conditions on a study sample of goshawk home ranges could take decades. Of course, interim monitoring and evaluating the effectiveness of implementation (the specific forest treatments) on moving the existing forest conditions toward the desired conditions and on

increasing the abundance of goshawk prey species should be undertaken. Such monitoring (versus a testing program focused on goshawks) could be achieved at greatly reduced costs because much smaller areas would be needed. Whatever approach is taken, a sound experimental design is required to evaluate implementation. Some potential problems in assessing the effectiveness of implementation are the needs for replications, risks of incorrectly assigning causal inferences due to ecological complexity and interactions within an ecosystem framework, and risks of spatial and temporal autocorrelations within the data (Mellina and Hinch 1995). Considerable economic costs would also be associated with testing the SWGS in sufficiently large landscapes. Because of these difficulties, combined with the improved likelihood that the broad-based ecosystem approach of the SWGS will successfully sustain goshawks, and because implementation initiates the restoration of management-altered forest habitats and ecosystems (see below), we suggest that immediate implementation in broad landscapes is a better option than the long wait for experimental tests of the SWGS's effectiveness. During implementation, however, we advocate monitoring programs that track the habitats and populations of goshawk and their prey, not necessarily within a testing framework, but as integral parts of an adaptive management program (McDonnell et al. 1997, Murry and Marmorek 2003). The SWGS was based on the habitat relationships of many plants and animals, an understanding of the autecology and synecology of the forest vegetation, and on knowledge of vegetation treatments to create the desired forest conditions. Do we know that this approach is appropriate or that the desired conditions are correct and sustainable (Long and Smith 2000)? Some degree of uncertainty exists regarding these questions; however, we do know that past management fundamentally altered forest ecosystems and that active management in many cases is needed to restore the ecosystems.

## ADDED BENEFITS OF IMPLEMENTATION

Reynolds et al. (1992) identified a number of added benefits from implementing the SWGS. A main benefit is restoration of forest ecosystems. Implementing of the SWGS benefits many plants and animals of southwestern forests by restoring tree densities, structures, and patterns similar to those occurring pre-settlement (circa 1850; Fig. 4). Throughout much of interior of western North America, tree densities in dry conifer forests (e.g., ponderosa pine and mixed-conifer forests) have greatly increased since the initiation of fire exclusion in the early 1900s (Cooper 1960, Weaver 1961, Covington and Moore 1994, Graham et al. 2004). In pre-settlement times, frequent surface fires maintained open forest conditions by cleaning the forest floor and killing small trees (Weaver 1943, Graham et al. 2004). In addition, timber harvests and associated treatments tended to homogenize forest composition, structure, and pattern (Nyland 2002). Thus, forests have become increasingly dense and less diverse. These changes increased inter-plant competition for moisture and nutrients, resulting in decreased tree vigor, increased tree disease and insect epidemics, and increased frequency of lethal wildfires (Weaver 1943, Fellin 1979, Williams and Marsden 1982, Anderson et al. 1987, Swetnam and Lynch 1989, Covington and Moore 1994, Graham 2003, Graham et al. 2004).

The desired forest conditions described in the SWGS resembled the historical conditions of southwestern ponderosa pine forests described by Pearson (1950) and White (1985). These similarities suggest that implementing the SWGS would move forests towards restoration of pre-settlement conditions (e.g., pre-timber harvest and fire exclusion) (Long and Smith 2000). For example, the SWGS restores old structures (large live trees, snags, and logs), maintains groups of trees with interlocking crowns, promotes the grass/forb/shrub layer, and minimizes the risk of lethal wildfires by reducing surface and ladder fuels (Reynolds et al. 1992, Graham 2003,

Graham et al. 2004). In addition, by favoring lower stand densities of seral species (e.g., ponderosa pine) the strategy reduces the likelihood of disease and insect epidemics (Schmid and Mata 1992, Harvey et al. 1999). These conditions also are similar to those suggested as being desirable in the Healthy Forests Initiative and Healthy Forests Restoration Act (USDA Forest Service 2004).

The SWGS has been described as single-species management (Beier and Maschinski 2003). However, the SWGS is a multi-species strategy because it was based on the habitats and ecological relationships of many plant and animals in the goshawk food web (Reynolds et al. 1992, Long and Smith 2000). Further, except for the nest areas and especially the inner portions of the PFA, goshawk habitat needs contributed relatively little to the desired landscape. Thus, the SWGS shifts the focus from single-species and stand-level management to vegetation management for food webs in large landscapes (Reynolds et al. 1992, Long and Smith 2000). The SWGS utilized the concept of desired forest conditions. Advantages of this concept include the recognition that long time periods may be required to attain the desired conditions, allows variable management actions depending on existing conditions, calls attention to native disturbance regimes and how these operated at multiple temporal and spatial scales, and focuses on resources that are left after treatment rather than on what resources are removed (Reynolds et al. 1992, Haynes et al. 1996, Graham et al. 1999, Franklin et al. 2002).

## SUMMARY

The strategy for conserving goshawks in the southwestern United States described desired forest landscapes intended to sustain the habitats of both goshawks and their prey (Reynolds et al. 1992). The approach and procedures developed in this conservation strategy are readily adapted to other locations and forests. However, the specific desired conditions for other forests are likely



to be different because the kinds of prey available as well as the composition, structure, pattern, and dynamics of the vegetation often differs among forests. The approach we present identifies goshawk habitats (nest and feeding) and habitats (nest and feeding) of important goshawk prey occurring in a forest (Fig. 1). Goshawk habitats were summarized in the SWGS, as were the habitats and life histories of 14 important goshawk prey species, a procedure readily applied to other forests and locales. Moreover, we described a procedure for reducing a full list of species eaten by goshawks to a manageable suite of important prey which is also applicable for other settings. The information assembled for the goshawk and its prey should be integrated with the ecological dynamics (e.g., regeneration, development, succession, and disturbances) of the vegetation in a focal forest type and we provided suggestions as to how this integration can be accomplished (Fig. 1). Depending on the current forest conditions (we provide suggestions on how they can be determined), management actions may be as simple as doing nothing, to actively managing (e.g., regenerating, thinning, prescribed burning) forests to develop and maintain goshawk and prey habitats. While we believe that the approach used in the SWGS for identifying and developing desired habitats for goshawks is sound, economically feasible, and, due to its diversity of components, robust to failure to sustain goshawks, we also realize that forest management is fraught with uncertainties and that managing goshawk and prey habitats is a long-term proposition. What is needed is an in-depth analysis of implementation projects as they come on line to make preliminary judgments about what works, what does not, and how success should be measured.

## ACKNOWLEDGEMENTS

We wish to thank our original coauthors who were instrumental in developing the SWGS. We also thank the many forest managers, scientists, and stakeholders who interacted with us over

the past 15 yr. Their many challenges caused us to frequently reexamine of our approach and desired forest conditions in the SWGS. In our minds, these reexaminations reaffirmed the appropriateness and value of our approach and its consequence. We thank T. Jain, K. Graham, D. Lund, and two anonymous reviewers of various drafts of the manuscript, and C. Erickson for help formatting the manuscript.

#### LITERATURE CITED

- Agee, J. K. 1993. Fire ecology of Pacific Northwest forests. Island Press, Washington, DC.
- Andersen, D. E., S. DeStefano, M. I. Goldstein, K. Titus, C. Crocker-Bedford, J. J. Keane, R. G. Anthony, and R. N. Rosenfield. 2004. The status of Northern Goshawks in the western United States. Technical Review. 04-1, The Wildlife Society, Bethesda, MA.
- Anderson, L., C. E. Carlson, and R. H. Wakimoto. 1987. Forest fire frequency and western spruce budworm outbreaks in western Montana. *Forest Ecology and Management* 22:251-260.
- Andersson, M., and R. Å. Norberg. 1981. Evolution of reversed sexual size dimorphism and role partitioning among predatory birds, with a size scaling of flight performance. *Biological Journal of the Linnean Society* 15:105–130.
- Baldwin, S. P., and S. C. Kendeigh. 1938. Variations in weights of birds. *Auk* 55:416–467.
- Bassett, R. L., D. A. Boyce, Jr., M. H. Reiser, R. T. Graham, and R. T. Reynolds. 1994. Influence of site quality and stand density on goshawk habitat in southwestern forests. *Studies in Avian Biology* 16:41–45.
- Bednarek, W., W. Hausdorf, V. Jörissen, E. Schulte, and H. Wegener. 1975. Über die Auswirkungen der chemischen Umweltbelastung auf Greifvögel in zwei Probeflächen Westfalens. *Journal für Ornithologie* 116:181–194.

- Beier, P., and J. E. Drennen. 1997. Forest structure and prey abundance in foraging areas of Northern Goshawks. *Ecological Applications* 7:564–571.
- Beier, P., and J. Maschinski. 2003. Threatened, endangered, and sensitive species. Pp. 306–327 in P. Friedrici (editor). *Ecological restoration of southwestern ponderosa pine forests*. Island Press, Inc., Washington, DC.
- Belovsky, G. E. 1987. Extinction models and mammalian persistence. Pp. 35–57 in M. E. Soulé (editor). *Variable populations for conservation*. Cambridge University Press, Cambridge, UK.
- Bielefeldt, J., R. N. Rosenfield, and J. M. Papp . 1992. Unfounded assumptions about diet of the Cooper's Hawk. *Condor* 94:427-436.
- Boal, C. W., D. E. Andersen, and P. L. Kennedy. 2003. Home range and residency of Northern Goshawks breeding in Minnesota. *Condor* 105:811–816.
- Boal, C. W., and R. W. Mannan. 1994. Northern Goshawk diets in ponderosa pine forests on the Kaibab Plateau. *Studies in Avian Biology* 16:97–102.
- Bond, R. M. 1940. A goshawk nest in the upper sonoran life-zone. *Condor* 42:100–103.
- Bosakowski, T., D. G. Smith, and R. Speiser. 1992. Niche overlap of two sympatric-nesting hawks *Accipiter* spp. in the New Jersey-New York Highlands. *Ecography* 15:358–372.
- Bosakowski, T., B. McCullough, F. J. Lapsansky, and M. E. Vaughn. 1999. Northern Goshawks nesting on a private industrial forest in western Washington. *Journal of Raptor Research* 33:240–244.
- Bosakowski, T., and D. G. Smith. 1992. Comparative diets of sympatric nesting raptors in the eastern deciduous forest biome. *Canadian Journal of Zoology* 70:984–992.
- Bradley, A. F., N. V. Noste, and W. C. Fischer. 1992. Fire ecology of forests and woodlands in

Utah. USDA Forest Service General Technical Report INT-287. USDA Forest Service, Intermountain Research Station, Ogden, UT.

- Braun, C. E., J. H. Enderson, Y. B. Linhart, C. D. Marti, and M. R. Fuller. 1996. Northern Goshawk and forest management in the southwestern United States. The Wildlife Society Technical Review 96-2.
- Bright-Smith, D. J., and R. W. Mannan. 1994. Habitat use by breeding male Northern Goshawks in northern Arizona. *Studies in Avian Biology* 16:58–65.
- Collins, C. T., and R. A. Bradley. 1971. Analysis of body weights of spring migrants in southern California: part II. *Western Bird Bander* 46:48–51.
- Cooper, C. F. 1960. Changes in vegetation, structure, and growth of southwestern pine forests since white settlement. *Ecological Monographs* 30:129–164.
- Cooper, S. V., K. E. Neiman, and D. W. Roberts. 1991. Forest habitat types of northern Idaho: A second approximation. USDA Forest Service. General Technical Report INT 236 USDA Forest Service, Intermountain Research Station, Ogden, UT.
- Covington, W. W., and M. M. Moore. 1994. Postsettlement changes in natural fire regimes and forest structure: Ecological restoration of old-growth ponderosa pine forests. Pp. 153–181 *in* R. N. Sampson, and D. L. Adams (editors). *Assessing forest ecosystem health in the Inland West*. The Haworth Press Inc, New York, NY.
- Cramp, S., and K. E. L. Simmons. 1980. *Birds of the western Palearctic*. Vol. 2. Hawks to bustards. Oxford University Press, Oxford, UK.
- Crocker-Bedford, D. C. 1990. Goshawk reproduction and forest management. *Wildlife Society Bulletin* 18:262–269.
- Daubenmire, R., and R. B. Daubenmire. 1968. Forest vegetation of eastern Washington and

northern Idaho. Washington Agriculture Experiment Station Technical Bulletin 60.

Washington State University, Pullman, WA.

Daw, S. K., and S. DeStefano. 2001. Forest characteristics of Northern Goshawk nest stands and post-fledging areas in Oregon. *Journal of Wildlife Management* 65:59-65.

DellaSala, D. A., J. R. Strittholt, R. F. Noss, and D. M. Olson. 1996 . A critical role for core reserves in managing Inland Northwest landscapes for natural resources and biodiversity. *Wildlife Society Bulletin* 24:209–221.

Dodd, N. L., J. S. States, and S. S. Rosenstock. 2003. Tassel-eared squirrel population, habitat condition, and dietary relationships in north-central Arizona. *Journal of Wildlife Management* 67:622–633.

Doyle, F. I., and J. M. N. Smith. 1994. Population responses of Northern Goshawks to the 10-year cycle in numbers of snowshoe hares. *Studies in Avian Biology* 16:122–129.

Drennan, J. E., and P. Beier. 2003. Forest structure and prey abundance in winter habitat of Northern Goshawks. *Journal of Wildlife Management* 67:177–185.

Dunning, J. B. 1984. Body weights of 686 species of North American birds. *Western Bird Bander Association Monograph* No. 1.

Everett, R. L., and J. F. Lehmkuhl. 1996. An emphasis-use approach to conserving biodiversity. *Wildlife Society Bulletin* 24:192–199.

Everett, R. L., and J. F. Lehmkuhl. 1997. A forum for presenting alternative viewpoints on the role of reserves in conservation biology? A reply to Noss (1996). *Wildlife Society Bulletin* 25:575–577.

Fellin, D. G. 1979. A review of some interactions between harvesting, residue management, fire, and forest insects and diseases. Pp. 335–415 *in* Environmental consequences of timber

harvesting in Rocky Mountain coniferous forests. USDA Forest Service General Technical Report INT-90. USDA Forest Service, Intermountain Forest and Range Experiment Station, Ogden, UT.

- Franklin, J. F., T. A. Spies, R. Van Pelt, A. B. Carey, D. A. Thornburgh, D. Rae Berg, D. B. Lindenmayer, M. E. Harmon, W. S. Keeton, D. C. Shaw, K. Bible, and J. Chen. 2002. Disturbances and structural development of natural forest ecosystems with silvicultural implications, using Douglas-fir forests as an example. *Forest Ecology and Management* 155:399-423.
- Fulé, P. Z., W. W. Covington, and M. M. Moore . 1997. Determining reference conditions for ecosystem management of Southwestern ponderosa pine forests. *Ecological Applications* 7:895-908.
- Good, R. E. 1998. Factors affecting the relative use of Northern Goshawk (*Accipiter gentilis*) kill areas in southcentral Wyoming. M.S. thesis, University of Wyoming, Laramie, WY.
- Graham, R. T. (technical editor). 2003. Hayman fire case study. USDA Forest Service. General Technical Report RMRS-GTR-114. USDA Forest Service, Rocky Mountain Research Station, Ogden, UT.
- Graham, R. T., S. McCaffrey, and T. B. Jain (technical editors). 2004. Science basis for changing forest structure to modify wildfire behavior and severity. USDA Forest Service General Technical Report RMRS-GTR-120. USDA Forest Service, Rocky Mountain Research Station, Ogden, UT.
- Graham, R. T., R. L. Rodriguez, K. M. Paulin, R. L. Player, A. P. Heap, and R. Williams. 1999. The Northern Goshawk in Utah: habitat assessment and management recommendations. USDA Forest Service General Technical Report RMRS-GTR-22. USDA Forest Service,

Rocky Mountain Research Station, Ogden, UT.

Grønnesby, S., and T. Nygård. 2000. Using time-lapse video monitoring to study prey selection by breeding Goshawks *Accipiter gentilis* in Central Norway. *Ornis Fennica* 77:117–129.

Guldin, J. M., D. Cawrse, R. Graham, M. Hemstrom, L. Joyce, S. Kessler, R. McNair, G.

Peterson, C. Shaw, P. Stine, M. Twery, and J. Walter. 2003. Science consistency reviews: a primer for application. USDA Forest Service FS-771. USDA Forest Service, Washington, DC.

Haig, I. T., K. P. Davis, and R. H. Weidman. 1941. Natural regeneration in the western white pine type. USDA Technical Bulletin No. 767. Washington, DC.

Hall, E. R. 1981. The mammals of North America. John Wiley & Sons, New York.

Hann, W. J., J. L. Jones, M. G. Karl, P. F. Hessburg, R. E. Keane, D. G. Long, J. P. Menakis, C.

H. McNicoll, S. G. Leonard, R. A. Gravenmier, and B. G. Smith. 1997. Landscape dynamics of the Basin. Pp. 338–1055 in T. M. Quigley, and S. J. Arbelbide (technical editors). An assessment of ecosystem components in the Interior Columbia Basin and portions of the Klamath and Great Basins: Volume II. USDA Forest Service General Technical Report PNW-405. USDA Forest Service, Pacific Northwest Research Station, Portland, OR.

Hargis, C. D., C. McCarthy, and R. D. Perloff. 1994. Home ranges and habitats of Northern Goshawks in eastern California. *Studies in Avian Biology* 16:66–74.

Hartman, F. A. 1955. Heart weight in birds. *Condor* 57:221–238.

Harvey, A. E., R. T. Graham, and G. I. McDonald. 1999. Tree species composition change-Forest soil organism interaction: potential affects on nutrient cycling and conservation processes in interior forests. Pp.137–145 in R. Meurisse, W. G. Ypsllantis, and C.

Seybold (technical editors). USDA Forest Service General Technical Report PNW-461.

USDA Forest Service, Pacific Northwest Research Station, Portland, OR.

Haynes, R. W., R. T. Graham, and T. M. Quigley. 1996. A framework for ecosystem management in the interior Columbia Basin and portions of the Klamath and Great Basins. USDA Forest Service and USDI Bureau of Land Management General Technical Report PNW-374. USDA Forest Service, Pacific Northwest Research Station, Portland, OR.

Herron, G. B., C. A. Mortimore, and M. S. Rawlings. 1985. Nevada raptors: their biology and management. Biological Bulletin No. 8, Nevada Department of Wildlife, Reno, NV.

Kaufmann, M. R., C. M. Regan, and P. M. Brown. 2000. Heterogeneity in ponderosa pine/Douglas-fir forests: age and size structure in unlogged and logged landscapes of central Colorado. Canadian Journal of Forest Research 30:698–711.

Keane, J. J. 1999. Ecology of the Northern Goshawk in the Sierra Nevada, California. Ph.D. dissertation, University of California, Davis, CA.

Kennedy, P. L. 1989. The nesting ecology of Cooper's Hawks and Northern Goshawks in the Jemez Mountains, NM. Final Report 35. USDA Forest Service, Santa Fe National Forest. Santa Fe, NM

Kennedy, P. L. 1990. Home ranges of Northern Goshawks nesting in north-central New Mexico. P. 259 in P. R. Klausman, and N. S. Smith (editors). Managing wildlife in the Southwest symposium. Arizona Chapter of the Wildlife Society, Phoenix, AZ.

Kennedy, P. L. 1991. Reproductive strategies of Northern Goshawks and Cooper's Hawks in north-central New Mexico. Ph.D. dissertation, Utah State University, Logan, UT.

Kennedy, P. L. 1998. Evaluating Northern Goshawk (*Accipiter gentilis atricapillus*) population



status: a reply to Smallwood and Crocker-Bedford. *Journal of Raptor Research* 32:336–342.

Kenward, R., and P. Widén. 1989. Do goshawks *Accipiter gentilis* need forests? Some conservation lessons from radio tracking. Pp. 561–567 in B.-U. Meyburd, and R. D. Chancellor (editors). *Raptors in the Modern World*. World Working Group on Birds of Prey and Owls, London, UK.

Kenward, R. E. 1979. Winter predation by Goshawks in lowland Britain. *British Bird* 72:64–73.

Kenward, R. E. 1982. Goshawk hunting behaviour, and range size as a function of food and habitat availability. *Journal of Animal Ecology* 51:69–80.

Kenward, R. E., V. Mareström, and M. Karlborn. 1981. Goshawk winter ecology in Swedish pheasant habitats. *Journal of Wildlife Management* 45:397–408.

Lewis, S. B., M. R. Fullon, and L. Titus. 2004. A comparison of 3 methods for assessing raptor diet during the breeding year. *Wildlife Society Bulletin* 32:373–385.

Lindén, H., and M. Wikman. 1980. Kanahaukan poikuekoon vaihtelusta suhteessa metsäkanalintujen runsauteen. [Brood size of the Goshawk in relation to tetraonid densities] . *Suomen Riista* 27:63–69.

Long, J. N., and F. W. Smith. 2000. Restructuring the forest, Goshawks and the restoration of southwestern ponderosa pine. *Journal of Forestry* 98:25–30.

MacCracken, J. G. 1996. Forest health in the inland Northwest: maintaining the focus. *Wildlife Society Bulletin* 24:325–329.

Martin, T. E. 1987. Food as a limit on breeding birds: a life-history perspective. *Annual Review of Ecology and Systematics* 18:453–487.

McDonnell, M. J., S. T. A. Pickett, P. Groffman, P. Bohlen, R. V. Pouyat, W. C. Zipperer, R. W.

- Parmelee, M. M. Carreiro, and K. Medley. 1997. Ecosystem processes along an urban-to-rural gradient. *Urban Ecosystems* 1:21–36.
- McGowan, J. D. 1975. Distribution, density and productivity of Goshawks in interior Alaska. Final Report Federal Aid in Wildlife Restoration, Project Report W17-3, 4, 5, 6. Juneau, AK.
- Melián, C. J., and J. Bascompte. 2002. Food web structure and habitat loss. *Ecology Letters* 5:37–46.
- Mellina, E., and S. G. Hinch. 1995. Overview of large-scale ecological experimental designs and recommendations for the British Columbia watershed restoration program. Watershed Restoration Project Report No. 1, Watershed Restoration Program, Ministry of Environment, Lands, and Parks, British Columbia, Canada. 28pp.
- Morin, P. J. 1981. Predatory salamanders reverse the outcome of competition among three species of anuran tadpoles. *Science* 212:1284–1286.
- Murray, C., and D. Marmorek. 2003. Adaptive management and ecological restoration. Pp. 417–428 in P. Friederici (editor). *Ecological restoration of southwestern ponderosa pine forests*. Island Press, Washington, DC.
- Noss, R. F. 1996. On attacking a caricature of reserves: response to Everett and Lehmkuhl. *Wildlife Society Bulletin* 24:777–779.
- Nyland, R. D. 2002. *Silviculture: concepts and applications*: 2nd Edition. McGraw-Hill, New York, NY.
- Oliver, C. D., and Larson, B. C. 1990. *Forest stand dynamics*. McGraw-Hill, New York, NY..
- O'Brien, R. A. 2002. *Arizona's Forest Resources, 1999*. USDA Forest Service Resource Bulletin RMRS-RB-2. USDA Forest Service, Rocky Mountain Research Station, Ogden, UT.

- Pearson, G. A. 1950. Management of ponderosa pine in the southwest. USDA Forest Service Agricultural Monograph No. 6. U.S. Government Printing Office, Washington, DC.
- Peck, J. 2000. Seeing the forest through the eyes of a hawk: an evaluation of recent efforts to protect Northern Goshawk populations in southwestern forests. *Natural Resources Journal* 40:125–156.
- Reich, R. M., S. M. Joy, and R. T. Reynolds. 2004. Predicting the location of Northern Goshawk nests: modeling the spatial dependency between nest locations and forest structure. *Ecological Modelling* 176:109–133.
- Reinhardt, E., and N. L. Crookston (technical editors). 2003. The fire and fuels extension to the forest vegetation simulator. USDA Forest Service General Technical Report RMRS-GTR-116. USDA Forest Service, Rocky Mountain Research Station, Ogden, UT.
- Reynolds, R. T. 1972. Sexual dimorphism in accipiter hawks: a new hypothesis. *Condor* 74:191-197.
- Reynolds, R. T. 1983. Management of western coniferous forest habitat for nesting accipiter hawks. USDA Forest Service General Technical Report RM-102. USDA Forest Service, Rocky Mountain Research Station, Ft. Collins, CO.
- Reynolds, R. T., E. C. Meslow, and H. M. Wight. 1982. Nesting habitat of coexisting Accipiter in Oregon. *Journal of Wildlife Management* 46:124-138.
- Reynolds, R. T., R. T. Graham, M. H. Reiser, R. L. Bassett, P. L. Kennedy, D. A. Boyce, Jr., G. Goodwin, R. Smith, and E. L. Fisher. 1992. Management recommendations for the Northern Goshawk in the southwestern United States. USDA Forest Service General Technical Report RM-217. USDA Forest Service, Rocky Mountain Research Station, Ft. Collins, CO.

- Reynolds, R. T., S. M. Joy, and D. G. Leslie. 1994. Nest productivity, fidelity, and spacing of Northern Goshawks in Arizona. *Studies in Avian Biology* 16:106–113.
- Reynolds, R. T., and E. C. Meslow. 1984. Partitioning of food and niche characteristics of coexisting *Accipiter* during breeding. *Auk* 101:761–779.
- Reynolds, R. T., J. D. Wiens, and S. R. Salafsky. In press. A review and evaluation of factors limiting Northern Goshawk populations. *Studies in Avian Biology*.
- Reynolds, R. T., and S. M. Joy. In press. Demography of Northern Goshawks in northern Arizona, 1991–1996. *Studies in Avian Biology*.
- Salafsky, S. R. 2004. Covariation between prey abundance and Northern Goshawk reproduction on the Kaibab Plateau, Arizona. M.S. thesis, Colorado State University, Ft. Collins, CO.
- Schmid, J. M., and S. A. Mata. 1992. Stand density and mountain pine beetle-caused tree mortality in ponderosa stands. USDA Forest Service General Technical Report RM-GTR-275. USDA Forest Service, Rocky Mountain Research Station, Ft. Collins, CO.
- Schmidt, K. M., J. P. Menakis, C. C. Hardy, W. J. Hann, and D. L. Bunnell. 2002. Development of coarse-scale spatial data for wildland fire and fuel management. USDA Forest Service General Technical Report RM-GTR-87. USDA Forest Service, Rocky Mountain Research Station, Ft. Collins, CO.
- Schnell, J. H. 1958. Nesting behavior and food habits of Goshawks in the Sierra Nevada of California. *Condor* 60:377–403.
- Selås, V. 1997. Influence of prey availability on re-establishment of goshawk *Accipiter gentilis* nesting territories. *Ornis Fennica* 74:113–120.
- Snyder, N. F., and J. W. Wiley. 1976. Sexual size dimorphism in hawks and owls of North America. *Ornithological Monographs* No. 20.

Sollien, A. 1979. Bestandsutviklingen hos hønschauk *Accipiter gentilis*, i Norge de siste 100 år.

[Goshawk population in Norway over the past 100 years.]. *Vår Fuglefauna* 2:95–106.

Sonsthagen, S. A. 2002. Year-round habitat, movement, and gene flow of Northern Goshawks breeding in Utah. M.S. thesis. Brigham Young University, Provo, UT.

Spies, T. A., and J. F. Franklin. 1996. The diversity and maintenance of old-growth forests. Pp. 296–314 in R. C. Szaro, and D. W. Johnson (editors). *Biodiversity in managed landscapes: theory and practice*. Oxford Press, New York, NY.

Squires, J. R., G. D. Hayward, and J. F. Gore. 1998. The role of sensitive species in avian conservation. Pp. 155–176 in J. M. Marzluff, and R. Sallabanks (editors). *Avian conservation: research and management*. Island Press, Washington, DC.

Squires, J. R., and R. T. Reynolds. 1997. Northern Goshawk (*Accipiter gentilis*). In A. Poole, and F. Gill (editors). *The Birds of North America*, # 298. The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologists' Union, Washington, DC.

Squires, J. R., and L. F. Ruggiero. 1995. Winter movements of adult Northern Goshawks that nested in southcentral Wyoming. *Journal of Raptor Research* 29:5–9.

Stephens, R. M. 2001. Migration, habitat use, and diet of Northern Goshawks that winter in the Uinta Mountains, Utah. M. S. thesis, University of Wyoming, Laramie, WY.

Storer, R. W. 1966. Sexual dimorphism and food habits in three North American *Accipiters*. *Auk* 83:423–436.

Sulkava, S., K. Huhtala, and R. Tornberg. 1994. Regulation of goshawk *Accipiter gentilis* breeding in western Finland over the last 30 years. Pp. 67–76 in B.-U. Meyburg, and R. D. Chancellor (editors). *Raptor conservation today. Proceedings of the IV World Conference on Birds of Prey and Owls*. Pica Press, London, UK.

- Suzuki, D. 2003. A look at world parks. *Science* 301:1289.
- Swem, T., and M. Adams. 1992. A Northern Goshawk nest in the tundra biome. *Journal of Raptor Research* 26:102.
- Swetnam, T. W., and A. M. Lynch. 1989. A tree-ring reconstruction of western spruce budworm history in the southern Rocky Mountains. *Forest Science* 35:962–986.
- Temeles, E. J. 1985. Sexual size dimorphism of bird-eating hawks: the effect of prey vulnerability. *American Naturalist* 125:485–499.
- Thomas, J. W., R. G. Anderson, C. Maser, and E. L. Bull. 1979. Snags. Pp. 60–77 in J. W. Thomas (editor). *Wildlife habitats in managed forest: the Blue Mountains of Oregon and Washington*. USDA Forest Service Agricultural Handbook No. 553. U.S. Government Printing Office, Washington, DC.
- Tornberg, R., and A. Colpaert. 2001. Survival, ranging, habitat choice and diet of the Northern Goshawk *Accipiter gentilis* during winter in Northern Finland. *Ibis* 143:41–50.
- USDA Forest Service. 1996. Record of decision for amending forest plans for Arizona and New Mexico. USDA Forest Service, Southwestern Region Albuquerque, NM.
- USDA Forest Service. 2004. The healthy forests initiative and healthy forests restoration act: interim field guide. USDA Forest Service and USDI Bureau of Land Management FS-799. Washington, DC.
- USGS National Gap Analysis Program. 1995. State gap analysis data. Utah Cooperative Fish and Wildlife Research Unit. Utah State University, Logan, UT.
- VanHorne, B. 1983. Density as a misleading indicator of habitat quality. *Journal of Wildlife Management* 47:893–901.
- Verner, J., and A. S. Boss. 1980. California wildlife and their habitats: western Sierra Nevada.

USDA Forest Service Pacific Southwest Forest and Range Experiment Station General  
Technical Report PSW-37. USDA Forest Service, Pacific Southwest Forest and Range  
Experiment Station, Berkley, CA.

- Weaver, H. 1943. Fire as an ecological and silvicultural factor in the ponderosa pine region of the  
pacific slope. *Journal of Forestry* 41:7–14.
- Weaver, H. 1961. Ecological changes in the ponderosa pine forest of Cedar Valley in southern  
Washington. *Ecology* 42:416–420.
- White, A. S. 1985. Presettlement regeneration patterns in a southwestern ponderosa pine stand.  
*Ecology* 66:589–594.
- White, C. M., G. D. Lloyd, and G. L. Richards. 1965. Goshawk nesting in the upper Sonoran in  
Colorado and Utah. *Condor* 67:269.
- White, G. C., and Garrott, R. A. 1990. Analysis of wildlife radio-tracking data. Academic Press,  
San Diego, CA.
- Widén, P. 1985. Breeding and movements of goshawks in boreal forests in Sweden. *Holarctic  
Ecology* 8:273-279.
- Widén, P. 1987. Goshawk predation during winter, spring and summer in a boreal forest area of  
central Sweden. *Holarctic Ecology* 10:104–109.
- Widén, P. 1989. The hunting habitats of goshawks *Accipiter gentilis* in boreal forests of central  
Sweden. *Ibis* 131:205–213.
- Widén, P. 1997. How, and why, is the Goshawk (*Accipiter gentilis*) affected by modern forest  
management in Fennoscandia? *Journal of Raptor Research* 31:107–113.
- Williams, R. E., and M. A. Marsden. 1982. Modeling probability of root disease center  
occurrence in northern Idaho forests. *Canadian Journal of Forest Research* 12:876–882.

- Woodbridge, B., and P. J. Detrich. 1994. Territory occupancy and habitat patch size of Northern Goshawks in the southern Cascades of California. *Studies in Avian Biology* 16:83–87.
- Wykoff, W. R., and R. A. Monserud. 1988. Representing site quality in increment models: A comparison of methods. Pp. 184–191 *in* A. R. Ek, S. R. Shifley, and T. E. Burk (editors). Forest growth modeling and prediction. USDA Forest Service, General Technical Report NC-120. USDA Forest Service, North Central Research Station, St. Paul, MN.
- Younk, J. V., and M. J. Bechard. 1994a. Breeding ecology of the Northern Goshawk in high-elevation aspen forests of northern Nevada. *Studies in Avian Biology* 16:119–121.
- Younk, J. V., and M. J. Bechard. 1994b. Effect of gold mining activity on Northern Goshawks breeding in Nevada's Independence and Bull Run Mountains. Final Report, Raptor Research Center, Boise State University, Boise, ID.



Table 1. An example for identifying a suite of important goshawk prey, including the numbers and percent frequency of individuals by species, and a frequency and biomass ranking of each species in diets of breeding goshawks in Oregon (29 species, 227 individuals; Reynolds and Meslow 1984).

SPECIES	Number <sup>a</sup>	Percent	Frequency	Biomass <sup>b</sup>
			Rank	Rank
Steller's Jay ( <i>Cyanocitta stelleri</i> )	29	12.8	1	4
Snowshoe hare ( <i>Lepus americanus</i> )	24	10.6	2	1
American Robin ( <i>Turdus migratorius</i> )	20	8.8	3	12
Golden-mantled ground squirrel ( <i>Citellus lateralis</i> )	17	7.4	4	5
Northern flying squirrel ( <i>Glaucomys sabrinus</i> )	15	6.6	5	7
Red-shafted Flicker ( <i>Colaptes auratus</i> )	15	6.6	5	10
Douglas' squirrel ( <i>Tamiasciurus douglasi</i> )	13	5.7	7	6
Mountain Quail ( <i>Oreortyx pictus</i> )	10	4.4	8	8
Mourning Dove ( <i>Zenaida macroura</i> )	7	3.1	9	16
Chipmunk spp. ( <i>Eutamias sp.</i> )	7	3.1	9	17
Blue Grouse ( <i>Dendragapus obscurus</i> )	5	2.2	11	2
Gray Squirrel ( <i>Sciurus grisesus</i> )	5	2.2	11	3

Gray Jay ( <i>Perisoreus canadensis</i> )	5	2.2	11	19
---	---	-----	----	----

-----Greater than 4 individuals/species threshold<sup>c</sup>-----

Belding's ground squirrel ( <i>Citellus beldingi</i> )	4	1.8	14	15
--	---	-----	----	----

Varied Thrush ( <i>Ixoreus naevius</i> )	4	1.8	14	20
--	---	-----	----	----

-----Greater than 3 individuals/species threshold<sup>c</sup>-----

Mountain cottontail rabbit ( <i>Sylvilagus nuttalli</i> )	3	1.3	16	11
---	---	-----	----	----

Townsend's chipmunk ( <i>Eutamias townsendi</i> )	3	1.3	16	23
---	---	-----	----	----

-----Greater than 2 individuals/species threshold<sup>c</sup>-----

Mallard ( <i>Anas platyrhynchos</i> )	2	0.9	18	9
---------------------------------------	---	-----	----	---

Ruffed Grouse ( <i>Bonasa umbellus</i> )	2	0.9	18	14
--	---	-----	----	----

Townsend's ground squirrel ( <i>Citellus townsendi</i> )	2	0.9	18	24
--	---	-----	----	----

European Starling ( <i>Sturnus vulgaris</i> )	2	0.9	18	27
---	---	-----	----	----

Hairy Woodpecker ( <i>Picoides villosus</i> )	2	0.9	18	30
---	---	-----	----	----

Williamson's Sapsucker ( <i>Sphyrapicus thyroideus</i> )	2	0.9	18	32
--	---	-----	----	----

Western Tanager ( <i>Piranga ludoviciana</i> )	2	0.9	18	34
--	---	-----	----	----

Finch spp. ( <i>Carpodacus sp.</i> )	2	0.9	18	36
Dark-eyed Junco ( <i>Junco hyemalis</i> )	2	0.9	18	39
Great horned Owl ( <i>Bubo virginianus</i> )	1	0.4	27	13
Cooper's Hawk ( <i>Accipiter cooperi</i> )	1	0.4	27	18
Bushy-tailed wood rat ( <i>Neotoma cinerea</i> )	1	0.4	27	21
Pileated Woodpecker ( <i>Dryocopus pileatus</i> )	1	0.4	27	22
Wood rat spp. ( <i>Neotoma sp.</i> )	1	0.4	27	25
Dusty-footed wood rat ( <i>Neotoma fuscipes</i> )	1	0.4	27	26
Black-billed Magpie ( <i>Pica pica</i> )	1	0.4	27	28
Western Screech-Owl ( <i>Otus kennicottii</i> )	1	0.4	27	29
Lewis' Woodpecker ( <i>Melanerpes lewis</i> )	1	0.4	27	31
Red-winged Blackbird ( <i>Agelaius phoeniceus</i> )	1	0.4	27	33
Red-naped Sapsucker ( <i>Sphyrapicus nuchalis</i> )	1	0.4	27	35
Black-headed Grosbeak ( <i>Pheucticus melanocephalus</i> )	1	0.4	27	37
Least chipmunk ( <i>Eutamias minimus</i> )	1	0.4	27	38

---

<sup>a</sup> After Reynolds and Meslow (1984).

<sup>b</sup> Biomass = number of individuals of a species in diet x mass of the species determined from the literature and museum collections (Reynolds and Meslow 1984; also see Baldwin and Kendeigh 1938, Hartman 1955, Collins and Bradley 1971, Dunning 1984, Bosakowski and Smith 1992) .

<sup>c</sup> The thresholds define three possible suites of important prey, with minimums of 4, 3, and 2 individual items per species. If the threshold of 4 individuals per species were used, the suite would contain 15 important prey, with some contributing as little as 1.8 % of items.

## FIGURE LEGENDS

FIGURE 1. Essential components and two levels of synthesis of goshawk habitats, prey habitats, and the composition, structure, and pattern of forests used to identify mixes of desired habitats in the southwestern goshawk conservation strategy (Reynolds et al. 1992).

FIGURE 2. The desired groups of trees with interlocking crowns surrounded by openings in Southwestern ponderosa pine forests.

FIGURE 3. Aerial photo showing a 1994 implementation of the southwestern goshawk conservation strategy (Reynolds et al. 1992) adjacent to seed tree harvests and a natural area in ponderosa pine forest in Arizona.

FIGURE 4. Historical mix of groups of different aged ponderosa pine trees on the Fort Valley Experimental Forest, Flagstaff Arizona (from Pearson 1950; see White 1985). This and other information (see text) provided references for supporting the desired sizes and mix of vegetation structural stages that could likely be sustained in Southwestern ponderosa pine forests (see Reynolds et al. 1992).




Fig. 1

Fig 2






Fig 3






Fig 4